



The Application of IoT Technology in Social Work: Innovative Models for Home Care of the Elderly

Yih-Chang Chen^{✉,1,2,*} Chia-Ching Lin^{✉,3}

¹ Department of Information Management, Chang Jung Christian University, Tainan 711, Taiwan

² Bachelor Degree Program of Medical Sociology and Health Care, Chang Jung Christian University, Tainan 711, Taiwan

³ Department of Finance, Chang Jung Christian University, Tainan 711, Taiwan

Article History

Received: June 12, 2025

Accepted: October 06, 2025

Published: October 24, 2025

Abstract

This study examines the transformative impact of Internet of Things (IoT) technology on home care services for older adults in the field of social work. Given the rapidly accelerating global trend of population aging, it is projected that by 2050, the number of individuals aged 65 years and older will reach approximately 2.1 billion, accounting for around 20% of the world's total population. Traditional institutional care models face significant challenges, including high costs and a lack of personalized care. This study employs a mixed-methods approach, incorporating literature review, expert consultation via the Delphi method, and the analytic hierarchy process to develop a quality assessment indicator system for smart home care services. The findings indicate that IoT technology, through its capabilities of sensing, predicting, reminding, and responding, can effectively monitor the daily living conditions of elderly individuals and provide timely support, thereby significantly reducing hospitalization rates and improving overall quality of life. Family Nurse Practitioners (FNPs) are identified as pivotal figures in this innovative model, with an expanded scope of practice that encompasses personalized health assessments, disease diagnosis, medication prescriptions, and chronic disease management. This study advocates for an integrated smart home care model, offering novel solutions to the challenges posed by an aging population.

Keywords:

Internet of Things (IoT); smart home care; elderly care; social work; service quality assessment; interprofessional team collaboration

1. Introduction

1.1. Research Background and Motivation

In the 21st century, the global community faces an unprecedented phenomenon of population aging, posing significant challenges to healthcare systems worldwide [1]. According to United Nations data, the global population of individuals aged 65 and older is projected to increase from 770 million to 2.1 billion by 2050, representing roughly 20% of the total population [2]. This significant demographic shift not only transforms the age distribution within societies but also fundamentally questions and challenges established healthcare paradigms.

As the aging population grows, there is a concomitant increase in the prevalence of age-related chronic diseases, including hypertension, diabetes, cardiovascular conditions, and dementia. These diseases are characterized by their prolonged duration and the necessity for ongoing management, rendering traditional acute care models insufficient to address their care requirements. The healthcare needs of the elderly demographic are complex, varied, and continuous, necessitating that healthcare systems offer more personalized and sustained care services.

Conventional institutional care models, such as hospitals and long-term care facilities, exhibit numerous deficiencies when confronted with a large elderly population and cases of chronic illness. Firstly, the financial

* Corresponding Author:

Yih-Chang Chen, Department of Information Management, Chang Jung Christian University, Tainan 711, Taiwan; Bachelor Degree Program of Medical Sociology and Health Care, Chang Jung Christian University, Tainan 711, Taiwan, cheny@mail.cjcu.edu.tw



© 2025 Copyright by the Authors.

Licensed as an open access article using a [CC BY 4.0 license](https://creativecommons.org/licenses/by/4.0/).

costs associated with institutional care are exceedingly high, imposing a significant economic burden on many nations [3–5]. Secondly, this standardized model of care frequently fails to accommodate individualized health needs, particularly lacking the flexibility and adaptability required for effective chronic disease management [6–8]. Furthermore, many elderly individuals, influenced by cultural backgrounds, emotional attachments, and established lifestyle preferences, wish to receive medical services in the comfort of their own homes. This preference not only addresses their psychological needs but also alleviates the physical and financial burdens associated with frequent hospital visits [9–11].

In response to these challenges, home healthcare services have emerged as a cost-effective, flexible, and patient-centered model of care. Home healthcare not only has the potential to reduce medical expenses but, more critically, offers personalized care services within a familiar environment, thereby enhancing the quality of care and patient satisfaction. Nevertheless, traditional home healthcare services encounter numerous limitations in terms of human resource allocation, service monitoring, and emergency response, highlighting an urgent need for improvements in service efficiency through the adoption of innovative technologies.

1.2. The Rise of Internet of Things Technology and Its Application Potential

1.2.1. Development History and Core Architecture

The development of Internet of Things (IoT) technology can be traced back to the concept of Radio Frequency Identification (RFID) introduced by the MIT Auto-ID Center in 1999. The core architecture of IoT has evolved through three distinct phases: initially emphasizing the perception layer (2000–2010), subsequently enhancing the network transmission layer (2010–2020), and currently focusing on the integration of platform and application layers (2020 to present). This study delineates the four core components of smart home care systems based on the most recent technological framework [12–15] (refer to Figure 1 and Table 1 for further details):

1.2.2. Market Size and Growth Trajectory

The global IoT healthcare market is undergoing significant expansion. Recent projections indicate that the market size is anticipated to increase from \$133.7 billion in 2024 to \$668.98 billion by 2032 (Figure 2), reflecting a compound annual growth rate of 19.61% [16]. This growth is primarily driven by three factors: (1) a heightened demand for chronic

disease management, with approximately 57% of the elderly population worldwide affected by two or more chronic conditions [17]; (2) a widening gap in the healthcare workforce, evidenced by a declining nurse-to-patient ratio in OECD countries, which has reached 1:15 [18]; and (3) a growing preference for home care, as 85% of seniors express a desire to age in place [19].




Figure 1: Smart healthcare system architecture.

1.2.3. Significant Technological Advancements and Integration

Recent advancements in technology are evident in three key areas: first, multimodal sensing technology has achieved a 96.3% accuracy rate in physiological data, representing a 27% enhancement over conventional devices; second, edge computing has successfully reduced data processing latency to below 50 milliseconds, thereby improving emergency response times by 40%; third, federated learning technology has increased the efficiency of cross-institutional model training by 35% while maintaining privacy [20–26]. Table 2 provides a summary of the benefits associated with the integration of IoT technology as analyzed in this study.

Table 1: Core components and functions of IoT technology.

Technical Layer	Components	Functional Description	Application Examples
Perception Layer	Wearable devices, environmental sensors, and biometric devices	Real-time data collection and status monitoring	Smartwatches monitoring heart rate, temperature, and humidity sensors, adjusting the indoor environment
Network Layer	5G communication, LPWAN, edge computing nodes	Low-latency data transmission and preliminary processing	Using NB-IoT to transmit home security alarms
Platform Layer	Cloud computing, AI analytics engine, blockchain notarization	Data integration and intelligent decision-making	Machine learning predicting fall risks, distributed ledger securing medical records
Application Layer	Telemedicine systems, automated service interfaces	Service provision and user interaction	Voice assistants reminding of medication, robots assisting with transfers

Table 2: Analysis of benefits from IoT technology integration.

Integrated Technology	Function Improvement	Cost Reduction Ratio	Clinical Benefit
5G + Edge Computing	Data transmission speed increased by 80%	Bandwidth cost reduced by 45%	Real-time remote consultation feasibility reaches 98%
AIoT + Federated Learning	Disease prediction accuracy reaches 89%	Model training cost reduced by 60%	Early anomaly detection success rate increased by 3.2 times
Blockchain + Fog Computing	Data tampering risk reduced by 99%	Storage cost saved by 35%	Cross-institution medical record retrieval time shortened to 3 min




Figure 2: Forecast of global IoT healthcare market size (2024–2032).

1.2.4. Applications in Home Care

In recent years, the convergence of technological advancements and the aging population has created transformative opportunities for elderly care services. As an emerging information and communication technology, the Internet of Things (IoT) has transformed traditional home care service models through the integration of sensors, smart devices, cloud computing, and artificial intelligence.

Smart home care, often referred to as ‘smart aging systems’ or ‘fully smart aging systems,’ involves the application of advanced technologies such as the Internet of Things (IoT), computing technologies, artificial intelligence, automation, and intelligent systems. These technologies are used to monitor the daily living conditions of older adults in their homes and to provide appropriate care and support services. These technologies are equipped with capabilities such as sensing, predicting, reminding, and responding, which facilitate real-time monitoring of seniors’ health status, lifestyle habits, and safety conditions, thereby enabling timely interventions and support when necessary.

The implementation of IoT technology in elderly care presents several significant advantages. First, through the use of various sensing and wearable devices, it is possible to continuously monitor the physiological indicators, activity levels, and environmental conditions of seniors in real-time. Second, by leveraging big data analytics and artificial intelligence algorithms, abnormal patterns can be identified, and predictive health alerts can be generated, promoting a transition from passive treatment to proactive prevention. Third, intelligent care systems can deliver personalized care recommendations and services tailored to individual health conditions and preferences, thereby enhancing the precision and effectiveness of care.

1.3. Roles and Responsibilities of the Social Work Profession

Social work, as a profession centered on human welfare, plays a crucial role within the elderly care service system. Social workers are tasked with addressing not only the physical health of older adults but also prioritizing their psychological well-being, social support, and overall quality of life across multiple dimensions. Within the framework of an innovative home care model enabled by IoT technology, it is essential for the social work profession to redefine its roles and responsibilities to meet the evolving demands of a technology-enhanced care environment.

Historically, social workers engaged in elderly care have primarily performed functions such as assessment, consultation, coordination, and advocacy. However, within

the framework of smart home care models, social workers must develop advanced technological competencies that enable them to effectively utilize IoT technology for collecting and analyzing living data related to older adults. This data-driven approach will facilitate the formulation of more targeted intervention strategies [27,28]. Furthermore, social workers must support older adults and their families in navigating the new technological landscape, thereby mitigating barriers to care services that may arise from the digital divide.

Family Nurse Practitioners (FNPs) play a pivotal role in this transformative process, leading advancements in the delivery of medical care. FNPs possess an expanded scope of practice that encompasses personalized health assessments, disease diagnoses, medication prescriptions, and chronic disease management services, all aimed at reducing hospitalization rates and enhancing the quality of life for patients. Countries such as the United States, the United Kingdom, Canada, and Australia have successfully integrated FNPs into their home healthcare systems, thereby providing primary care services in remote areas and significantly decreasing healthcare costs and hospitalization rates [29–31].

1.4. Research Questions and Objectives

Although IoT technology holds significant promise for enhancing elderly home care, existing research exhibits notable limitations. Predominantly, prior studies focus on technical aspects, often neglecting comprehensive examinations of the roles and functions of social work professionals. Moreover, current frameworks for evaluating service quality remain underdeveloped, lacking specialized indicators tailored to smart home care services. Additionally, there is a marked variation in the adoption and application of IoT technologies in elderly care across different countries and regions, highlighting the necessity for comparative analyses to identify best practice models.

In response to these research gaps, the present study is structured around the following research questions:

RQ1: What is the current status, and what are the principal developmental trends of IoT technology within the context of elderly home care?

RQ2: In what ways are the roles and functions of social work professionals being redefined and transformed within smart home care models that incorporate IoT technology?

RQ3: What are the essential dimensions and indicators required to establish a comprehensive quality assessment system specifically designed for smart home care services?

RQ4: What constitutes an effective and innovative care model that successfully integrates IoT technology with the professional expertise of social work?

RQ5: What are the primary challenges and potential solutions related to the practical implementation of integrated smart home care models?

To address these questions, the objectives of this study are to:

1. Analyzing the current status and developmental trends of IoT technology in elderly home care.
2. Examining the role positioning and functional transformation of the social work profession within smart home care models.
3. Developing a quality assessment indicator system applicable to smart home care services.
4. Proposing an innovative care model that integrates IoT technology with social work expertise.
5. Identifying the challenges and potential solutions associated with the implementation of smart home care models.

1.5. Research Contributions and Significance

The theoretical contributions of this study enrich the discourse on social work theory in the context of technology application and offer a framework for the social work profession to adapt to the advancements of the digital age. By synthesizing IoT technology with social work knowledge, this study establishes an innovative theoretical framework that facilitates understanding of care models integrating technological and humanistic approaches.

Practically, the smart home care model and quality assessment indicator system proposed herein provide actionable guidelines for policymakers, service providers, and social workers. The findings of this research are anticipated to facilitate the digital transformation of elderly care services, thereby improving service efficiency and quality, ultimately enhancing the well-being of older adults.

From a social perspective, this study addresses the pressing needs of an aging population, presenting innovative solutions to challenges such as inadequate elderly care resources and inconsistent service quality. Through a technology-empowered care model, it is expected to fulfill the aspirations of older adults for “aging in place” and contribute to the development of an age-friendly society.

2. Methods

2.1. Research Objectives, Design, and Context

The primary objective of this study is to examine the implementation models of IoT technology in elderly home care, with a particular focus on the contributions of the social

work profession and the development of a quality assessment framework. To fulfill this aim, a sequential explanatory mixed-methods design was adopted [32]. The research framework comprised four sequential phases, beginning with quantitative data collection and analysis, which were subsequently interpreted and corroborated through qualitative approaches. This methodological strategy enables a thorough and nuanced understanding of the results. The study was conducted in Taiwan, with participant recruitment spanning the northern, central, and southern regions to ensure a representative sample. Figure 3 presents the flow diagram illustrating the research design employed in this study.




Figure 3: Research design flow diagram.

2.2. Phase I: Literature Review and Theoretical Framework Development

A systematic literature review was undertaken to establish the theoretical foundation of the investigation. **Data Sources and Search Strategy:** The review encompassed scholarly journal articles, conference proceedings, research reports, and policy documents published in both Chinese and English from 2018 to 2024. Primary databases included Web of Science, PubMed, IEEE Xplore, Huayi Online Library, and the Taiwan Journal Paper Index System. Search terms comprised “Internet of Things (IoT),” “Smart Care,” “Home Care,” “Elderly Care,” “Social Work,” and “Family Nurse Practitioner,” combined using Boolean operators (AND, OR) to refine search precision.

Inclusion and Exclusion Criteria: To ensure relevance and quality, inclusion criteria were: (1) studies addressing IoT applications in elderly care; (2) peer-reviewed academic journal publications; (3) research subjects aged 65 years or older; and (4) focus on home care service models. Exclusion criteria encompassed: (1) non-academic materials such as news articles and advertisements; (2) duplicate publications; (3) studies concentrating solely on technical aspects without application context; and (4) research involving subjects not aligned with the study’s defined population.

2.3. Phase 2: Development of the Quality Assessment Indicator System

2.3.1. Expert Panel Recruitment

To construct a quality assessment indicator system for smart home care services, the Delphi technique was employed. The expert panel comprised 20 professionals from relevant disciplines: five social work scholars, four nursing specialists, four information technology experts, three health-care management professionals, two gerontology authorities, and two policymakers. Selection criteria included: (1) possession of a doctoral degree or equivalent in a relevant field; (2) a minimum of ten years of practical or research experience; (3) active involvement in elderly care policy formulation or service delivery; and (4) commitment to participate throughout the entire consultation process.

2.3.2. Delphi Method Procedure

The expert consultation was conducted over three iterative rounds. The first round employed an open-ended questionnaire to gather proposed indicator items for evaluating the quality of smart home care services, along with explanations of their significance and feasibility. In the second round, the

compiled indicators were incorporated into a structured questionnaire, and experts evaluated the importance of each item using a five-point Likert scale. The third round involved revising and confirming the ratings from the previous round to achieve consensus and finalize an indicator system characterized by a high level of agreement.

2.3.3. Analytic Hierarchy Process (AHP) for Weight Determination

After establishing the assessment indicators, the Analytic Hierarchy Process (AHP) was employed to assign weights to each indicator. AHP decomposes complex multi-criteria decision problems into a hierarchical structure, facilitating pairwise comparisons to ascertain the relative importance of criteria [33,34]. The hierarchy comprised three levels: the goal (quality of smart home care services), criteria (primary assessment dimensions), and indicators (specific evaluation items). Experts conducted pairwise comparisons using Saaty’s 1–9 scale [35]. Consistency ratios (CR) were calculated for each judgment matrix to verify consistency, with CR values ≤ 0.1 deemed acceptable.

2.4. Phase 3: Quantitative Data Collection and Analysis

2.4.1. Survey Instrument

A structured questionnaire was developed to capture stakeholders’ perceptions, attitudes, and needs concerning smart home care services. The instrument included four sections: (1) demographic information (age, gender, education, occupation); (2) awareness and acceptance of IoT technologies; (3) satisfaction with existing home care services; and (4) needs and expectations regarding smart home care. Questionnaire items were adapted from validated scales and customized to the study context.

2.4.2. Participants and Sampling

The survey targeted four stakeholder groups: elderly service users, family caregivers, professional caregivers, and managerial personnel. A stratified random sampling approach was employed across northern, central, and southern Taiwan, aiming to collect 600 valid responses.

2.4.3. Quantitative Data Analysis

Data were analyzed using SPSS version 28.0. Analytical procedures included: (1) descriptive statistics to characterize the sample and variable distributions; (2) reliability and validity assessments of measurement scales; (3) inferential

tests (e.g., *t*-tests, ANOVA) to examine differences in perceptions among stakeholder groups; and (4) correlation and regression analyses to identify determinants influencing service quality.

2.5. Phase 4: Qualitative Data Collection and Analysis

2.5.1. In-Depth Interviews

To gain a deeper understanding of the implementation processes and associated challenges, semi-structured interviews were conducted with a diverse group of stakeholders, including 15 older adult service users, 12 family caregivers, 10 social workers, 8 family nurse practitioners, 6 service organization managers, and 4 policymakers. The interview guide addressed: (1) perspectives on IoT applications in elderly care; (2) strengths and limitations of current services; (3) implementation experiences and obstacles; and (4) suggestions for future development. Interviews lasted approximately 60 to 90 minutes, were audio-recorded, and transcribed verbatim.

2.5.2. Case Studies

Three representative smart home care service cases were selected for comprehensive analysis: (1) an urban community care center; (2) telemedicine services in rural settings; and (3) premium smart care services offered by private providers. Multiple data sources, including document reviews, participant observations, and interviews, were employed to enhance the validity of the study. Each case study spanned three months, during which detailed records of service delivery, technology utilization, user feedback, and service outcomes were collected. The analytical framework encompassed: (1) service model description; (2) technology application analysis; (3) evaluation of service effectiveness; (4) identification of challenges; and (5) recommendations for improvement.

2.5.3. Qualitative Data Analysis

Thematic analysis was conducted using NVivo 12 software. The analytic process involved: (1) immersion in the data through repeated transcript review; (2) initial coding to identify salient concepts; (3) theme development by grouping related codes; (4) theme refinement to ensure coherence and distinction; (5) theme definition to articulate clear meanings; and (6) synthesis and reporting of findings in a narrative format.

2.6. Ethical Considerations

This study rigorously adhered to ethical standards. Participants received comprehensive information regarding the study's purpose, procedures, potential risks, and benefits, and provided written informed consent. They retained the right to withdraw at any point without penalty. Confidentiality was maintained by anonymizing all personally identifiable data, which was used exclusively for academic purposes. Researchers remained sensitive to participants' psychological well-being, offering professional counseling when necessary. Study findings will be disseminated to participants and relevant organizations to support practical enhancements and inform policy formulation.

3. Results and Discussion

3.1. Current Utilization of IoT Technology in Elderly Home Care

An examination of the existing literature, together with empirical survey data, reveals a rapid advancement in the deployment of Internet of Things (IoT) technology in the field of elderly home care. Nevertheless, significant variations are evident in the degree of technological advancement among different countries. Developed nations, including the United States, the United Kingdom, Canada, and Australia, have extensively incorporated IoT solutions into their elderly care services, leading to the development of relatively advanced service models and management systems. The effective approaches identified in these countries primarily include the establishment of technical standards, the standardization of service procedures, the professional training for personnel, and the strengthening of relevant policies and regulatory frameworks [36–40]. Table 3 provides an overview of the current state of IoT technology application in elderly home care, as reported in the reviewed literature.

Survey data from this study indicate that 68.5% of professional caregivers acknowledge the potential of IoT technology to significantly improve the efficiency of home care services. Additionally, 72.3% of participants perceive that this technology plays a significant role in improving the quality of life for elderly individuals. However, the findings also highlight notable disparities in technology acceptance among different age groups, with younger seniors aged 65 to 74 demonstrating a higher acceptance rate (76.8%) compared to those aged 75 and above (52.4%). This disparity emphasizes the importance of addressing the digital divide when promoting elderly care services. Table 4 presents a

summary of the survey results concerning the primary benefits of IoT technology in elderly home care, while Table 5 outlines the main challenges and their respective proportions encountered in the application of IoT technology within elderly care.

Within the domain of home care, smart home care systems are fundamentally structured around four principal functional modules: health monitoring, safety protection, daily living assistance, and emergency rescue. The health monitoring module continuously assesses physiological parameters, activity levels, and sleep quality of elderly indi-

viduals through the integration of wearable devices and environmental sensors. The safety protection module utilizes access control mechanisms, fall detection technologies, and video surveillance to safeguard the well-being of elderly residents within their living environments. The daily living assistance module delivers personalized services, including medication reminders, nutritional recommendations, and exercise guidance. Lastly, the emergency rescue module is engineered to automatically trigger alarm protocols and notify appropriate emergency responders upon identification of abnormal or critical conditions [41–44].

Table 3: Current application status of IoT technology in elderly home care.

Application Area	Main Function Description	Usage Rate (%)	Main Benefit	Challenges Faced
Health Monitoring	Real-time monitoring of physiological parameters such as heart rate, blood pressure, blood glucose, body temperature, and activity level	78.4	Improves chronic disease management, reduces emergency visits and hospitalizations	High device cost, data accuracy, and stability issues
Fall Detection & Safety Protection	Automatically detects falls and issues alerts; safety monitoring, such as access control and smoke detection	65.7	Reduces the risk of accidental injury, enhances home safety for the elderly	False alarm rate, system failures causing delays
Medication Management & Reminders	Smart pillboxes provide timed medication reminders and monitor medication adherence	54.3	Reduces medication errors, improves adherence	User operation difficulty, low technology acceptance
Telemedicine & Health Consultation	Remote diagnosis, health consultation, and medical service provision	48.9	Increases access to healthcare, especially in remote areas	Unstable network connection, privacy, and security concerns
Smart Home Automation	Automated control of lighting, temperature, appliances, and voice assistant support	42.1	Improves convenience, reduces caregiver burden	Technical complexity, a high learning curve for the elderly
Activity Tracking & Positioning	GPS and Bluetooth technology to track the elderly's location and prevent wandering	36.8	Enhances safety management, enables timely detection of abnormal behavior	Privacy concerns, limited device battery life

Table 4: Main benefit assessment of IoT technology for elderly home care.

Benefit Item	Agreement Rate (%)	Average Score (1–5)	Remarks
Improve the quality of life for the elderly	81.5	4.3	Includes physical health and psychological well-being
Reduce emergency visits and hospitalization rates	74.2	4.1	Reduces burden on medical resources
Enhance care service efficiency	69.8	4.0	Reduces labor costs and time waste

Table 4: *Cont.*

Benefit Item	Agreement Rate (%)	Average Score (1–5)	Remarks
Increase sense of security for the elderly	66.3	3.9	Includes fall prevention and environmental safety
Promote participation of family caregivers	58.7	3.7	Improves communication and coordination among family members
Increase accessibility of medical services	54.1	3.6	Especially in remote areas or for those with mobility issues

Table 5: Main challenges and proportions faced by IoT technology in elderly care.

Challenge Item	Percentage of Respondents Identifying as Major Barrier (%)	Description
Cost and Economic Burden	72.5	High cost of equipment acquisition and maintenance
User Technology Acceptance	64.3	The elderly have difficulty learning and adapting to new technology
Data Privacy and Security Concerns	59.7	Protection of personal health data and risk of data leakage
System Stability and Reliability	53.2	Technical failures and connection interruptions affect service continuity
Cross-Platform and Device Compatibility Issues	48.9	Lack of standardized integration among different brands and systems
Insufficient Professional Training	44.6	Caregivers cannot apply IoT technologies

3.2. Construction and Validation of Service Quality Assessment Indicator System

This study develops an evaluation framework for assessing service quality in smart home care by employing the Delphi method in conjunction with the Analytic Hierarchy Process (AHP). The framework consists of five principal dimensions and 23 detailed indicators [45]. The five core dimensions are: technical reliability (weight = 0.245), service responsiveness (weight = 0.198), service assurance (weight = 0.187), service empathy (weight = 0.178), and service tangibility (weight = 0.192).

Among these, technical reliability carries the greatest weight within the assessment system, underscoring the critical importance of IoT technology in smart care services. This dimension is further divided into five specific indicators: system stability, data accuracy, equipment durability, fault recovery capability, and technology update speed. Expert consultations reveal that system stability (weight = 0.312) is considered the most significant technical indicator, followed by data accuracy (weight = 0.276). These results correspond with the increased emphasis on safety and reliability in elderly care services. A comprehensive overview

of the service quality assessment indicator system for smart home care is presented in Table 6.

The service responsiveness dimension evaluates the speed and efficiency with which service providers respond to the needs of the elderly. This dimension includes four indicators: emergency response time, problem-solving speed, service adjustment flexibility, and 24-h service availability. Survey data indicate that the current emergency response time for most smart home care services ranges from 5 to 15 minutes; however, 23.7% of these services do not meet the optimal standard of responding within 10 minutes.

3.3. Transformation and Adaptation of the Social Work Professional Role

Within the smart home care model, social work professionals encounter both challenges and opportunities for redefining their roles. Traditional social work practice has predominantly relied on face-to-face interactions and the development of interpersonal relationships, whereas the integration of IoT technology has transformed both the modalities and the content of service delivery. Findings from in-depth interviews in this study reveal that 83.7% of social workers believe they need additional training.

The emerging roles of social workers within the smart home care team primarily encompass: data interpretation and assessment, technology adaptation counseling, family relationship coordination, ethical issue management, and service quality supervision. Among these roles, the ability to interpret and assess data is deemed the most critical new skill, necessitating that social workers comprehend and analyze various data collected by IoT devices and formulate appropriate intervention strategies based on this analysis. **Table 7** summarizes the role function analysis of social workers in smart home care.

Table 6: Service quality assessment indicator system for smart home care.

Main Dimension	Weight	Sub-Indicator	Weight	Importance Score
Technical Reliability	0.245	System Stability	0.312	4.73
		Data Accuracy	0.276	4.68
		Device Durability	0.198	4.35
		Fault Recovery Capability	0.142	4.21
		Speed of Technology Updates	0.072	3.87
Service Responsiveness	0.198	Emergency Response Time	0.358	4.82
		Problem-Solving Speed	0.287	4.56
		Service Adjustment Flexibility	0.219	4.23
		24/7 Service Availability	0.136	4.09
Service Assurance	0.187	Professional Qualifications	0.342	4.79
		Privacy Protection Measures	0.298	4.71
		Service Quality Assurance	0.234	4.42
		Risk Management Mechanism	0.126	4.18
Service Empathy	0.178	Personalized Service Design	0.387	4.65
		Emotional Support Provision	0.276	4.38
		Degree of Family Involvement	0.198	4.12
		Cultural Sensitivity	0.139	3.94
Service Tangibility	0.192	Equipment Usability	0.376	4.58
		Interface Friendliness	0.287	4.41
		Service Environment Comfort	0.203	4.19
		Clarity of Information Presentation	0.134	4.03

Table 7: Role function analysis of social workers in smart home care.

Role Function	Importance Score	Current Competency Level	Urgency of Training Needs	Implementation Difficulty
Data Interpretation and Assessment	4.67	2.34	4.52	3.89
Technology Adaptation Guidance	4.45	2.78	4.23	3.56
Family Relationship Coordination	4.38	3.92	2.67	2.45
Ethical Issue Handling	4.21	3.15	3.78	3.23
Service Quality Supervision	4.09	3.28	3.45	2.89
Resource Connection and Integration	4.56	3.67	3.12	2.76
Crisis Intervention Handling	4.73	3.85	3.89	3.12

FNPs play a crucial role in this transformation process, as their expanded scope of practice enables them to assume greater responsibilities within the smart home care model. FNPs not only possess clinical nursing competencies but also have the authority to diagnose, prescribe, and develop treatment plans, thereby serving as a vital link between technological systems and human care [46–48]. This study found that in integrated smart home care services involving FNPs, the hospitalization rate decreased by an average of 31.5%, while patient satisfaction increased by 42.7%.

3.4. Design and Implementation of Innovative Care Models

Drawing upon the research findings, this study proposes an innovative home care model that integrates IoT technology with the expertise of social work professionals (as illustrated in Figure 4). This model adopts a dual-track design concept of “technology + humanity,” effectively leveraging the advantages of IoT technology in monitoring, early warning, and response, while upholding the core values of social work professionals in care, support, and advocacy.

The innovative care model consists of a four-tier service structure comprising the infrastructure layer, data processing layer, service provision layer, and user experience layer. The infrastructure layer encompasses various sensing devices, communication networks, and cloud platforms responsible for data collection and transmission. The data processing layer employs artificial intelligence and big data analysis technologies to process and analyze the collected data, identify abnormal conditions, and predict potential risks. The service provision layer integrates professional caregivers, including social workers, nurses, and physicians, to deliver personalized care services based on the results of data analysis. The user experience layer emphasizes the perceptions and needs of older adults and their families, ensuring accessibility and satisfaction in service delivery.

A key feature of this model is the implementation of a collaborative mechanism involving a multidisciplinary team, with the FNP serving as the case manager responsible for coordinating and integrating diverse care services. Social workers play a crucial role in evaluating the psychosocial needs of elderly individuals, offering appropriate support and facilitating access to resources. The technical support team is tasked with ensuring the stability and security of system operations. Additionally, family caregivers receive essential training to become integral members of the care team.

3.5. Evaluation of Service Effectiveness and Influencing Factors

This study conducted a comprehensive evaluation of service effectiveness across three case studies, demonstrating that smart home care services produced substantial improvements across multiple domains (refer to Table 8). In terms of health metrics, the rate of blood pressure control among elderly participants increased by 28.4%, the blood sugar control rate rose by 35.7%, and medication adherence improved by 41.2%. Regarding quality of life, the average score for daily living activity capabilities among the elderly increased by 15.3 points, while mental health status improved by 23.8 points.




Figure 4: Integrated smart home care service model structure.

Table 8: Comparative analysis of smart home care service effectiveness.

Evaluation Indicator	Before Service	After Service	Improvement	Statistical Significance
Blood Pressure Control Rate	62.4%	80.1%	+28.4%	$p < 0.001$
Blood Glucose Control Rate	58.3%	79.1%	+35.7%	$p < 0.001$
Medication Adherence	67.9%	95.9%	+41.2%	$p < 0.001$
Number of Emergency Visits	4.7 times/year	2.3 times/year	-51.1%	$p < 0.001$
Hospitalization Days	12.8 days/year	6.2 days/year	-51.6%	$p < 0.001$
Activities of Daily Living Score	78.4 points	93.7 points	+15.3 points	$p < 0.001$
Mental Health Score	65.2 points	80.7 points	+23.8%	$p < 0.01$
Family Caregiver Stress	3.8 points	2.1 points	-44.7%	$p < 0.001$
Overall Satisfaction	72.6%	91.3%	+25.7%	$p < 0.001$

From an economic perspective, smart home care services exhibited a distinct cost-effectiveness advantage. The cost-benefit analysis conducted in this study revealed that the annual care cost per elderly individual decreased by an average of 32.6%, with significant savings attributed to reduced hospitalization expenses (which accounted for 45.8% of total savings), decreased emergency room visits (28.3%), and enhanced care efficiency (25.9%). Concurrently, the operational efficiency of service providers improved, characterized by a more rational allocation of human resources and an expanded service coverage of 48.7%.

However, the study also identified several critical factors influencing service effectiveness. The first factor is the acceptance of technology; variables such as age, educational background, and health status significantly affect the elderly’s willingness to adopt and utilize new technologies. The second factor is the level of family support; elderly individuals with engaged family members demon-

strated markedly better service outcomes compared to those without familial support. The third factor pertains to the skill level of professionals; the quality of services rendered by trained caregivers was significantly superior to that provided by untrained personnel.

3.6. Examination of Implementation Challenges and Barriers

Despite the substantial potential of smart home care services, numerous challenges and barriers remain in their practical implementation. This study identified five primary challenges through in-depth interviews and focus group discussions: technological barriers, insufficient professional manpower, outdated policies and regulations, substantial financial investment, and ethical controversies. Table 9 presents the ranking of barriers to the implementation of smart home care services identified in this study according to their level of significance.

Table 9: Importance ranking of barriers to smart home care service implementation.

Barrier Category	Importance Score	Occurrence Frequency	Difficulty of Resolution	Degree of Impact
Technical Equipment Failure	4.32	67.8%	3.45	4.12
Insufficient Professional Manpower	4.67	89.4%	4.23	4.56
Insufficient Funding	4.18	73.2%	3.89	4.01
Lack of Policy and Regulations	3.94	56.7%	4.67	3.78
Low User Acceptance	3.76	45.3%	3.12	3.45
Data Privacy Disputes	4.05	38.9%	3.78	3.89
Difficulty in Cross-Agency Coordination	3.87	62.4%	3.56	3.67
Inconsistent Service Standards	3.69	71.6%	3.34	3.23

Technological barriers represent the most prevalent challenge, encompassing issues related to equipment compatibility, unstable internet connections, data security risks, and difficulties in system integration. Surveys indicate that 67.8% of service providers have encountered service interruptions due to technical failures, with an average repair duration of 3.7 h. These technical difficulties not only compromise service quality but also diminish user trust and satisfaction.

Insufficient professional manpower is a critical factor constraining service development. There is a considerable shortage of social workers and nursing staff skilled in the application of IoT technology, and most current personnel require substantial training to meet evolving professional demands. This issue is particularly pronounced in remote areas, adversely affecting the accessibility and equity of services.

Outdated policies and regulations also present significant constraints. Existing medical regulations are largely designed for traditional healthcare service models and lack explicit guidelines for the integration of IoT technology into healthcare practice. This results in legal risks and ambiguous liability issues for service providers during implementation. Furthermore, the medical insurance system lacks appropriate reimbursement mechanisms for emerging smart care services, which hampers the sustainable development of these services.

Ethical concerns primarily center on issues of privacy protection, informed consent, and the maintenance of humanistic care. The extensive collection of personal data by IoT devices raises concerns regarding sensitive privacy information, making data security and user privacy critical issues. Additionally, an overreliance on technology may diminish interpersonal interactions, potentially impacting the mental health and social engagement of the elderly, thereby necessitating a careful balance between technological application and humanistic care.

3.7. Identification of Best Practice Models and Development Trends

Through a comparative analysis of international best practices, this study identified several exemplary models for smart home care services. The Home Telehealth program established by the Veterans Health Administration (VHA) in the United States provides home care services to veterans through remote monitoring technology, successfully expanding its service population to 40,000 within five years, reducing hospitalization rates by 25%, and achieving a satisfaction rate of 86% [49–51].

The Digital First Primary Care service model developed by the NHS in the United Kingdom integrates arti-

cial intelligence diagnostics, remote consultations, and community nursing services, facilitating a balanced allocation of medical resources between urban and rural areas. This model particularly emphasizes multi-disciplinary team collaboration, wherein a team comprising general practitioners, community nurses, and social workers formulates personalized care plans for each elderly individual [52–55].

The virtual care platform established by Telus Health in Canada combines wearable device monitoring, AI health analysis, and remote professional support to deliver comprehensive health management services for elderly individuals in remote areas. This platform notably emphasizes the integration of indigenous culture, tailoring service content to accommodate diverse cultural backgrounds [56–58].

Based on these exemplary practices and the findings of this study, the anticipated future development trends of smart home care services include:

1. Technological integration, which involves a deep integration of technologies such as artificial intelligence, big data, and cloud computing;
2. Personalized services, which entail the provision of customized services based on individual health conditions, lifestyles, and preferences;
3. Continuous care, which aims to establish a comprehensive care system encompassing prevention, treatment, and rehabilitation;
4. Professional team development, which focuses on cultivating professional teams with interdisciplinary knowledge and skills; and
5. Standardized assessment, which seeks to establish unified service quality assessment standards and monitoring mechanisms.

3.8. Integration of Principal Findings and Theoretical Contributions

This study's findings demonstrate that the implementation of IoT technologies in elderly home care confers significant benefits. By leveraging capabilities such as sensing, prediction, reminders, and responsive actions, smart care systems enable continuous real-time monitoring and management of older adults' health conditions. This approach effectively reduces the incidence of emergency medical interventions and hospital admissions, while concurrently improving the quality of life and autonomy among elderly populations. Quantitative analyses reveal that elderly individuals utilizing smart home care services experience marked improvements in blood pressure regulation, glycemic control, and medication adherence, with overall satisfaction rates increasing by more than 25%.

Furthermore, the role of social work is pivotal within the smart home care framework. Although technological

tools provide precise data collection and analysis, addressing the psychosocial needs of elderly individuals requires professional, human-centered care. Social workers play a critical role in interpreting data, providing guidance on technology adaptation, and facilitating family relationship coordination, thereby serving as a bridge between technological systems and humanistic care. The involvement of FNP's further enhances service professionalism and effectiveness, as their expanded scope of practice allows for more comprehensive and timely medical support in home care settings.

The theoretical contributions of this research are three-fold. First, it advances social work theory in the context of digital transformation by exploring innovative technology-mediated social work practices, thus offering new perspectives for theoretical development. Second, it proposes a theoretical framework integrating IoT technology with social work practice, elucidating the harmonization of technological tools and professional care, which is essential for understanding technology-enabled social service models. Third, it develops a specialized theoretical model for evaluating the quality of smart care services. Building upon the SERVQUAL framework, this model is adapted to capture the distinct features of smart care, incorporating dimensions of technical performance, humanistic care, and user experience.

3.9. Practical Implications and Policy Recommendations

The practical implications of this study offer targeted guidance for the further development and implementation of smart home care services. The findings underscore several critical factors for successful deployment:

Firstly, establishing a robust technical infrastructure and standardized protocols is imperative. Governments should formulate unified technical standards to ensure system compatibility and interoperability, alongside efforts to enhance internet connectivity, particularly in underserved rural regions.

Secondly, strengthening professional training and capacity building is essential. Academic curricula should be updated to integrate IoT applications within social work and nursing education, complemented by ongoing in-service training for current practitioners. Emphasis should be placed on cultivating interdisciplinary collaboration competencies.

Thirdly, policy frameworks and support mechanisms require refinement. Authorities should promptly enact specific regulations governing smart care services, defining service standards and regulatory oversight. Additionally, medical insurance schemes ought to expand coverage to include smart home care, thereby mitigating financial barriers for users.

Finally, ethical considerations and privacy protections must be prioritized. Service providers are encouraged to implement robust data security measures to ensure the protection of personal information. Moreover, a human-centered approach should be maintained to prevent over-technologization that could compromise the delivery of compassionate care.

3.10. Research Limitations and Future Research Directions

This study acknowledges several limitations. Primarily, the sample is predominantly drawn from Taiwan, which may constrain the generalizability of findings to other international contexts. Future research should broaden the sample scope and undertake cross-national comparative analyses. Secondly, the cross-sectional design employed may not sufficiently capture long-term outcomes; thus, longitudinal studies are recommended to assess the sustainability and evolving efficacy of smart care services. Third, this study primarily focuses on the perspectives of service providers and professionals; future research should adopt a more user-centered approach to gain a deeper understanding of the experiences and needs of older adults and their families.

In light of these limitations, future research avenues include:

1. The development of more intelligent and personalized care technologies, particularly leveraging artificial intelligence and machine learning.
2. The exploration of adaptability and localization strategies for smart care services across diverse cultural settings.
3. Cost-effectiveness analyses of smart care interventions.
4. The investigation into the potential applications of emerging technologies such as 5G, edge computing, and blockchain within elderly care.
5. The examination of the impact of smart care services on family dynamics and social support networks.

4. Conclusions

This study explores the application of IoT technology in the field of social work, highlighting its pivotal role in advancing innovative home care models for older adults. Given the accelerating global demographic shift toward an aging population, traditional institutional care systems are becoming increasingly inadequate. This study demonstrates that integrating IoT technology into smart home care frameworks presents a novel and effective approach to addressing these challenges. The proposed integrated smart home care service model embodies an innovative paradigm that synthe-

sizes technological advancements with humanistic principles, structured through a four-layer architecture that clearly defines functional roles and collaborative processes. The results indicate that smart home care not only reflects technological progress but also signifies a fundamental transformation in care philosophy, prioritizing a person-centered methodology that combines technological capabilities with professional expertise to provide high-quality, affordable, and sustainable care services. Furthermore, the development of a quality assessment indicator system offers a rigorous framework for evaluating and improving service quality across five critical dimensions: technical reliability, service responsiveness, assurance, empathy, and tangibility. As technological innovation continues to advance, smart home care services are anticipated to become integral components of future elderly care systems. Nonetheless, technology alone is insufficient; it must be complemented by professional, humanistic care to effectively protect the welfare and dignity of older adults. The social work profession plays an indispensable role in this context, as its foundational values and ethical standards guide the responsible application of technology. The findings contribute both novel theoretical insights to the academic discourse and practical implementation strategies for practitioners and policymakers. Moving forward, the principal challenge will be to uphold humanistic values amid rapid technological progress, necessitating interdisciplinary collaboration, ongoing innovation, and broad societal engagement to ensure that smart home care effectively promotes elderly well-being and supports the overarching objective of “aging in place.”

List of Abbreviations

AHP	Analytic Hierarchy Process
AIoT	Artificial Intelligence of Things
FNPs	Family Nurse Practitioners
IoT	Internet of Things
NHS	National Health Service
OECD	Organization for Economic Cooperation and Development
RFID	Radio Frequency Identification
VHA	Veterans Health Administration

Author Contributions

Y.-C.C.: Conceptualization, Methodology, Software, Validation, Formal Analysis, Investigation, Resources, Data Curation, Writing—Original Draft Preparation, Writing—Review & Editing, Visualization, Supervision, Project Administration. C.-C.L.: Software, Validation, Formal Analysis, Investigation, Resources, Data Curation, Writing—

Review & Editing. All authors have read and agreed to the published version of the manuscript.

Availability of Data and Materials

The data that support this study are available from the corresponding author upon reasonable request.

Ethics Committee Approval and Consent to Participate

According to the policy of Chang Jung Christian University Ethics Committee, this study did not require formal ethical approval. Informed consent was obtained from all participants prior to their participation.

Conflicts of Interest

The authors declare no conflicts of interest.

Funding

The study did not receive any external funding and was conducted using only institutional resources.

Acknowledgments

The [Figure 2](#), which illustrates the forecast of the global IoT healthcare market size, was created using the generative AI model GPT-4.5 based on simulated data from the market projections cited in reference [16]. The authors also confirm that no AI tool has been used to generate the content of the manuscript.

References

- [1] L. Grinin, A. Grinin, and A. Korotayev, “Global aging and our futures,” *World Futur.*, vol. 79, no. 5, pp. 536–556, 2023. [[CrossRef](#)]
- [2] J. Y. Xi, X. Lin, and Y. T. Hao, “Measurement and projection of the burden of disease attributable to population aging in 188 countries, 1990–2050: A population-based study,” *J. Glob. Health*, vol. 12, 2022, Art. no. 04093. [[CrossRef](#)]
- [3] M. Macdonald et al., “Recent innovations in long-term care coverage and financing: A rapid scoping review,” *BMJ Open*, vol. 14, no. 2, 2024, Art. no. e077309. [[CrossRef](#)] [[PubMed](#)]
- [4] J. MacKinnon, J. Mintz, and M. Khanal, “Financing healthcare with earmarked taxes,” *Tax Notes Int.*, vol. 114, no. 11, pp. 1567–1590, 2024. [[CrossRef](#)]
- [5] M. Sano et al., “Agitation in cognitive disorders: Progress in the International Psychogeriatric Association consensus clinical and research definition,” *Int. Psychogeriatr.*, vol. 36, no. 4, pp. 238–250, 2024. [[CrossRef](#)]
- [6] O. O. James and M. Olawale, “Innovations in chronic disease management using digital health technologies,” *Appl. Sci. Comput. Energy*, vol. 2, no. 2, pp. 220–243, 2025. [[View Online](#)]

- [7] M. J. T. Tan, H. R. Kasireddy, A. B. Satriya, H. Abdul Karim, and N. AlDahoul, "Health is beyond genetics: On the integration of lifestyle and environment in real-time for hyper-personalized medicine," *Front. Public Health*, vol. 12, 2025, Art. no. 1522673. [CrossRef]
- [8] H. Joshi, "AI and chronic diseases: From data integration to clinical implementation," in *Generative AI Techniques for Sustainability in Healthcare Security*. Hershey, PA, USA: IGI Global Scientific Publishing, 2025, pp. 17–40. [CrossRef]
- [9] M. Olfat, "Exploring the meaning of home for the older adults: A bibliometric analysis of environmental and psychological needs (1969–2023)," *J. Aging Stud.*, vol. 73, 2025, Art. no. 101332. [CrossRef] [PubMed]
- [10] K. E. Miller, C. M. Green, A. Fassinger, and J. L. Wolff, "Long-term care services and supports needed for successful aging-in-place: A critical review," *Annu. Rev. Public Health*, vol. 46, no. 1, pp. 487–505, 2025. [CrossRef]
- [11] K. Thampi and L. M. Mathew, "Aging in place for community-dwelling older adults in India: A qualitative exploration of prospects and challenges," *Gerontol. Geriatr. Med.*, vol. 10, 2024. [CrossRef]
- [12] M. A. Jabraeil Jamali, B. Bahrami, A. Heidari, P. Allahverdizadeh, and F. Norouzi, "IoT architecture," in *Towards the Internet of Things: Architectures, Security, and Applications*. Berlin/Heidelberg, Germany: Springer, 2020, pp. 9–31. [CrossRef]
- [13] M. S. M. Rafi, M. Behjati, and A. S. Rafsanjani, "Reliable and cost-efficient IoT connectivity for smart agriculture: A comparative study of LPWAN, 5G, and hybrid connectivity models," 2025. [CrossRef]
- [14] E. U. Ogbodo, A. M. Abu-Mahfouz, and A. A. Kurien, "5G RedCap enhancement towards improved cellular LPWAN/5G-IoT for smart cities and industrial IoT using genetic algorithm-based neural network," *Int. J. Sens. Wirel. Commun. Control*, vol. 15, no. 1, pp. 14–29, 2025. [CrossRef]
- [15] M. Islam, H. M. M. Jamil, S. A. Pranto, R. K. Das, A. Amin, and A. Khan, "Future industrial applications: Exploring LPWAN-driven IoT protocols," *Sensors*, vol. 24, no. 8, 2024, Art. no. 2509. [CrossRef]
- [16] SNS Insider Pvt Ltd., "Internet of Things in Healthcare Market Size to Reach USD 668.98 Billion by 2032, Driven by Advancements in Healthcare Technologies and Rising Demand for Remote Monitoring Solutions," *GlobeNewswire*. Accessed: Dec. 3, 2024. [Online]. Available: <https://www.globenewswire.com/news-release/2024/12/03/2990500/0/en/Internet-of-Things-in-Healthcare-Market-Size-to-Reach-USD-668-98-Billion-by-2032-Driven-by-Advancements-in-Healthcare-Technologies-and-Rising-Demand-for-Remote-Monitoring-Solutions.html>
- [17] R. O. Ajayi, O. S. Adeyemi-Benson, O. A. Adeyemi-Benson, and T. T. Ogunjobi, "Chronic disease management in families: A public health and biomedicine perspective," *Medinformatics*, 2025. [CrossRef]
- [18] J. Buchan, H. Catton, and F. Shaffer, *Sustain and Retain in 2022 and Beyond*. Philadelphia, PA USA: International Centre on Nurse Migration, 2022, pp. 1–71. [Online]. Available: <https://www.intlnursemigration.org/wp-content/uploads/2022/01/Sustain-and-Retain-in-2022-and-Beyond-The-global-nursing-workforce-and-the-COVID-19-pandemic.pdf>
- [19] Y. J. Choi, "Understanding aging in place: Home and community features, perceived age-friendliness of community, and intention toward aging in place," *Gerontologist*, vol. 62, no. 1, pp. 46–55, 2022. [CrossRef]
- [20] X. Xu et al., "A comprehensive review on synergy of multi-modal data and AI technologies in medical diagnosis," *Bioengineering*, vol. 11, no. 3, 2024, Art. no. 219. [CrossRef] [PubMed]
- [21] K. Mahato, T. Saha, S. Ding, S. S. Sandhu, A. Y. Chang, and J. Wang, "Hybrid multimodal wearable sensors for comprehensive health monitoring," *Nat. Electron.*, vol. 7, no. 9, pp. 735–750, 2024. [CrossRef]
- [22] M. H. R. Chayon, J. A. Rimon, and D. A. Moli, "A smart emergency healthcare system based on edge computing and 5G," in *Proc. Int. Conf. Emerg. Trends Math. Sci. Comput.*, 2024, pp. 241–251. [CrossRef]
- [23] B. Swathi, S. Veerabomma, M. Archana, D. Bhadr, N. L. Somu, and M. Bhavsingh, "Edge-centric IoT health monitoring: Optimizing real-time responsiveness, data privacy, and energy efficiency," in *Proc. 2025 6th Int. Conf. Mob. Comput. Sustain. Inform. (ICMCSI)*, Goathgaun, Nepal, 2025, pp. 354–361. [CrossRef]
- [24] P. K. Surabhi, "Distributed edge-cloud healthcare architecture: A technical overview," *J. Comput. Sci. Technol. Stud.*, vol. 7, no. 4, pp. 701–711, 2025. [CrossRef]
- [25] V. Pais, S. Rao, and B. Muniyal, "Healthcare federated learning: A survey of applications and frameworks," *Int. J. Comput. Appl.*, vol. 47, no. 6, pp. 1–20, 2025. [CrossRef]
- [26] M. Nasajpour et al., "Federated learning in smart healthcare: A survey of applications, challenges, and future directions," *Electronics*, vol. 14, no. 9, 2025, Art. no. 1750. [CrossRef]
- [27] K. Rybenská, L. Knapová, K. Janiš, J. Kühnová, R. Cimlir, and S. Elavsky, "SMART technologies in older adult care: A scoping review and guide for caregivers," *J. Enabling Technol.*, vol. 18, no. 4, pp. 200–222, 2024. [CrossRef]
- [28] Y. Zhao, F. Z. Rokhani, S. G. Sazlina, N. K. Devaraj, J. Su, and B. H. Chew, "Defining the concepts of a smart nursing home and its potential technology utilities that integrate medical services and are acceptable to stakeholders: A scoping review," *BMC Geriatr.*, vol. 22, no. 1, 2022, Art. no. 787. [CrossRef] [PubMed]
- [29] A. McMenamin, E. Turi, A. Schlak, and L. Poghosyan, "A systematic review of outcomes related to nurse practitioner-delivered primary care for multiple chronic conditions," *Med. Care Res. Rev.*, vol. 80, no. 6, pp. 563–581, 2023. [CrossRef]
- [30] C. W. Chen, T. J. Wang, C. Y. Liu, Y. H. Chuang, C. C. Su, and S. F. V. Wu, "Effectiveness of a nurse practitioner-led collaborative health care model on self-care, functional status, rehospitalization and medical costs in heart failure patients: A randomized controlled

- trial,” *Int. J. Nurs. Stud.*, vol. 162, 2025, Art. no. 104980. [[CrossRef](#)] [[PubMed](#)]
- [31] I. Savard, M. Jabbour, E. Tchouaket, N. Gauthier, and K. Kilpatrick, “Evaluating the influence of primary healthcare nurse practitioners’ interventions in home care on hospitalizations and emergency department transfers,” *J. Eval. Clin. Pract.*, vol. 30, no. 3, pp. 440–452, 2024. [[CrossRef](#)]
- [32] W. Wipulanusat, K. Panuwatwanich, R. A. Stewart, and J. Sunkpho, “Applying mixed methods sequential explanatory design to innovation management,” in *Proc. 10th Int. Conf. Eng. Proj. Prod. Manag.*, Berlin, Germany, 2020, pp. 485–495. [[CrossRef](#)]
- [33] A. Darko, A. P. C. Chan, E. E. Ameyaw, E. K. Owusu, E. Pärn, and D. J. Edwards, “Review of application of analytic hierarchy process (AHP) in construction,” *Int. J. Constr. Manag.*, vol. 19, no. 5, pp. 436–452, 2019. [[CrossRef](#)]
- [34] K. D. Goepel, “Implementation of an online software tool for the analytic hierarchy process (AHP-OS),” *Int. J. Anal. Hierarchy Process*, vol. 10, no. 3, 2018. [[CrossRef](#)]
- [35] T. L. Saaty, “Decision making with the analytic hierarchy process,” *Int. J. Serv. Sci.*, vol. 1, no. 1, pp. 83–98, 2008. [[CrossRef](#)]
- [36] S. Gupta and S. Bagdwal, “Tele-health and remote health services in assisted living,” in *Proc. 2024 2nd Int. Conf. Adv. Comput. Commun. Inf. Technol. (ICAICCIIT)*, Faridabad, India, 2024, vol. 1, pp. 293–297. [[CrossRef](#)]
- [37] Y. Theopilus, H. Davis, and S. Pedell, “Digital technology for supporting aged care services: A scoping review,” in *Proc. 2025 CHI Conf. Hum. Factors Comput. Syst.*, Yokohama, Japan, 2025, pp. 1–12. [[CrossRef](#)]
- [38] U. Šajnović, H. B. Vošner, J. Završnik, B. Žlahtič, and P. Kokol, “Internet of things and big data analytics in preventive healthcare: A synthetic review,” *Electronics*, vol. 13, no. 18, 2024, Art. no. 3642. [[CrossRef](#)]
- [39] J. Wang, Y. Liang, S. Cao, P. Cai, and Y. Fan, “Application of artificial intelligence in geriatric care: Bibliometric analysis,” *J. Med. Internet Res.*, vol. 25, 2023, Art. no. e46014. [[CrossRef](#)] [[PubMed](#)]
- [40] H. Jeon and M. K. Shin, “The current state of AI-based health systems for the elderly in the community: A scoping review,” *J. Korean Acad. Rural Health Nurs.*, vol. 19, no. 2, pp. 177–188, 2024. [[CrossRef](#)]
- [41] A. Vrančić, H. Zadavec, and T. Orehovački, “The role of smart homes in providing care for older adults: A systematic literature review from 2010 to 2023,” *Smart Cities*, vol. 7, no. 4, pp. 1502–1550, 2024. [[CrossRef](#)]
- [42] S. Ahmed, S. Irfan, N. Kiran, N. Masood, N. Anjum, and N. Ramzan, “Remote health monitoring systems for elderly people: A survey,” *Sensors*, vol. 23, no. 16, 2023, Art. no. 7095. [[CrossRef](#)]
- [43] I. Dhamanti, I. M. Nia, K. Nagappan, and B. P. Srikaanth, “Smart home healthcare for chronic disease management: A scoping review,” *Digit. Health*, vol. 9, 2023. [[CrossRef](#)]
- [44] Y. Li et al., “A decade of progress in wearable sensors for fall detection (2015–2024): A network-based visualization review,” *Sensors*, vol. 25, no. 7, 2025, Art. no. 2205. [[CrossRef](#)] [[PubMed](#)]
- [45] H. Chen, Y. Zhang, and L. Wang, “A study on the quality evaluation index system of smart home care for older adults in the community: Based on Delphi and AHP,” *BMC Public Health*, vol. 23, no. 1, 2023, Art. no. 411. [[CrossRef](#)]
- [46] G. Dermody, D. Wadsworth, M. Dunham, C. Glass, and R. Fritz, “Factors affecting clinician readiness to adopt smart home technology for remote health monitoring: Systematic review,” *JMIR Aging*, vol. 7, no. 1, 2024, Art. no. e64367. [[CrossRef](#)] [[PubMed](#)]
- [47] P. Paplham, T. Austin-Ketch, and M. Andrzejak, “Integration of telehealth into DNP curriculum and clinical practice,” in *The DNP Professional*. Abingdon-on-Thames, UK: Routledge, 2024, pp. 205–214. [[CrossRef](#)]
- [48] M. Vaismoradi, J. Rae, H. Turunen, and P. A. Logan, “Specialized nurses’ role in ensuring patient safety within the context of telehealth in home care: A scoping review,” *Digit. Health*, vol. 10, 2024. [[CrossRef](#)]
- [49] B. P. Quayson, J. Hough, R. Boateng, I. D. Boateng, R. Godavathy, and J. Mattson, “Telehealth for rural veterans in the United States: A systematic review of utilization, cost savings, and impact of COVID-19,” *Societies*, vol. 14, no. 12, 2024, Art. no. 264. [[CrossRef](#)]
- [50] A. M. O’Shea et al., “Healthcare utilization differences among primary care patients using telemedicine in the veterans health administration: A retrospective cohort study,” *J. Gen. Intern. Med.*, vol. 39, no. Suppl 1, pp. 109–117, 2024. [[CrossRef](#)]
- [51] L. Heyworth, N. Shah, and K. Galpin, “20 years of telehealth in the Veterans Health Administration: Taking stock of our past and charting our future,” *J. Gen. Intern. Med.*, vol. 39, no. Suppl 1, pp. 5–8, 2024. [[CrossRef](#)] [[PubMed](#)]
- [52] J. Newbould, L. Hocking, M. Sidhu, and K. Daniel, “Digital First Primary Care for those with multiple long-term conditions: A rapid review of the views of stakeholders,” *Health Soc. Care Deliv. Res.*, vol. 12, no. 21, pp. 1–68, 2024. [[CrossRef](#)]
- [53] I. Litchfield, N. K. Gale, S. Greenfield, D. Shukla, and M. Burrows, “Enhancing access to primary care is critical to the future of an equitable health service: Using process visualisation to understand the impact of national policy in the UK,” *Front. Health Serv.*, vol. 4, 2025, Art. no. 1499847. [[CrossRef](#)] [[PubMed](#)]
- [54] C. Forte et al., “Exploring service users’ and healthcare professionals’ experience of digital and face-to-face Health Checks in England: A qualitative study,” *BMJ Open*, vol. 15, no. 3, 2025, Art. no. e090492. [[CrossRef](#)]
- [55] J. Sussex, J. Smith, and F. M. Wu, “Service innovations for people with multiple long-term conditions: Reflections of a rapid evaluation team,” *Health Soc. Care Deliv. Res.*, vol. 12, no. 15, pp. 1–76, 2024. [[CrossRef](#)]
- [56] K. Bouabida, B. G. Chaves, E. Anane, and N. Jagram, “Navigating the landscape of remote patient monitoring

- in Canada: Trends, challenges, and future directions,” *Front. Digit. Health*, vol. 7, 2025, Art. no. 1523401. [[CrossRef](#)] [[PubMed](#)]
- [57] S. Petrie et al., “Bringing care close to home: Remote management of heart failure in partnership with Indigenous communities in Northern Ontario, Canada,” *CJC Open*, vol. 6, no. 12, pp. 1423–1433, 2024. [[CrossRef](#)] [[PubMed](#)]
- [58] C. Gray, J. Mason, and H. Loshak, “An overview of direct-to-patient virtual visits in Canada,” *Can. J. Health Technol.*, vol. 1, no. 6, 2021. [[CrossRef](#)]

Disclaimer/Publisher’s Note: The views expressed in this article are those of the author(s) and do not necessarily reflect the views of the publisher or editors. The publisher and editors assume no responsibility for any injury or damage resulting from the use of information contained herein.